

What are the harmful/adverse effects of multiple doses of COVID-19 mRNA vaccines?

<https://covid19evidencereviews.saskhealthauthority.ca/en/permalink/coviddoc439>

Document Type: Rapid Review
Review Code: EOC220103 RR
Question Submitted: January 20, 2022
Date Completed: February 4, 2022
Status: 3. Completed
Research Team: EOC

Key Findings:

Evidence to date of adverse events associated with mRNA COVID-19 vaccines are largely local reactions such as pain and swelling of the injection site, and systemic reactions of an allergic nature

Increased incidences of myocarditis and pericarditis have been observed, most often occurring in men under 30 years of age, and most often after the second dose of mRNA vaccine, with more occurrences following the Moderna COVID-19 vaccine versus Pfizer, however all cases appear to resolve

To date, there is no evidence indicating a negative effect on the immune system from the administration of multiple doses of mRNA COVID-19 vaccines

Category: Clinical Presentation
Healthcare Services

Subject: Infection Prevention and Control
Risk
Vaccines

Population: All

Clinical Setting: Primary care
Public Health

Priority Level: Level 3 Two weeks (14 days)

Cite As: Badea, A; Reeder, B; Groot, G; Miller, L; Mueller, M. What are the harmful/adverse effects of multiple doses of COVID-19 mRNA vaccines? 2022 Feb 04, Document no.: EOC220103 RR. In: COVID-19 Rapid Evidence Reviews [Internet]. SK: SK COVID Evidence Support Team, c2022. 10 p. (CEST rapid review report).

Related Documents:

Evidence Search Report: EOC220103 ESR
Table: EOC220103 RR Table

Documents



EOC220103 RR

 Read PDF

 Download PDF